

On the local unitary equivalence of states of multi-partite systems

M. Ziman¹, P. Štelmachovič¹, and V. Bužek^{1,2}

¹ *Research Center for Quantum Information, Slovak Academy of Sciences, Dúbravská cesta 9, 842 28 Bratislava, Slovakia*

² *Faculty of Informatics, Masaryk University, Botanická 68a, 602 00 Brno, Czech Republic*
(25 June 2001)

Two pure states of a multi-partite system are always related by a unitary transformation acting on the Hilbert space of the whole system. This transformation involves multi-partite transformations. On the other hand some quantum information protocols such as the quantum teleportation and quantum dense coding are based on equivalence of some classes of states of bi-partite systems under the action of local (one-particle) unitary operations. In this paper we address the question: “Under what conditions are the two states ϱ and σ , of a multi-partite system locally unitary equivalent?” We present a set of conditions which have to be satisfied in order that the two states are locally unitary equivalent. In addition, we study whether it is possible to prepare a state of a multi-qudit system, which is divided into two parts A and B , by unitary operations acting only on the systems A and B , separately.

I. INTRODUCTION

Quantum entanglement is the key ingredient in quantum information processing [1]. Bipartite entanglement is quite “well” understood by now, but the investigation of the multi-partite case is just beginning.

Several quantum information protocols such as quantum teleportation and quantum dense coding are based on the equivalence of some classes of states of bi-partite systems under the action of local (one-particle) unitary operations [1,2]. So the question of local unitary equivalence of multi-partite states of quantum systems is of importance. Recently several authors [4–6] have studied the so-called *polynomial invariants* of local unitaries. In the present paper we discuss the problem of local unitary equivalence. We present a set of necessary conditions which the two multi-partite states have to fulfill in order to declare them *locally unitary equivalent*.

The state of a quantum system is usually represented by a *density matrix*. We know that the elements of the density matrix depend on the choice of the basis in Hilbert space, $\mathcal{H}$, of the system. Physically the basis can be thought of as a *complete set of observables*. For example, in the case of spin-1/2 particle, the choice of the basis in Hilbert space corresponds to the direction of the Stern-Gerlach experimental setup [3].

A local unitary transformation corresponds to a change of a basis in each of the subsystems. Such unitary transformations are known as *passive* transformations, because there is no physical action performed on the quantum system itself. These transformations simply reflect the choice of our point of view rather than any specific manipulation of the physical system. It reflects the fact that two locally-unitary-equivalent states have the same matrix form, only the choice basis of subsystems is different. An *active* unitary transformation corresponds to the most general dynamical map of the states for isolated systems. It means that under the dynamics described by

local unitaries each of the subsystems evolves independently. Physically it corresponds to the absence of interactions between the subsystems. This property allows us to prepare from a given state of the multi-partite system some other state only by using *local* unitaries (here we assume that the subsystems can be separated by space-like intervals). From a mathematical point of view there is no difference between the active and passive unitary transformations.

In this paper we address the question: “Under what conditions are the two states ϱ and σ , of a multi-partite system locally unitary equivalent?” We present a set of conditions which have to be satisfied in order that the two states are locally unitary equivalent. In addition we study whether it is possible to prepare a state of a multi-qudit system which is divided into two parts A and B by unitaries acting only on the systems A and B , separately. In Section II we present examples of some invariants together with some applications for pure states. A criterion about the local unitary equivalence for mixed states is given in Section III. In Section IV we investigate the connection of bipartite and multi-partite entanglement via local unitaries for system of N qudits.

II. LOCAL UNITARY EQUIVALENCE

Let us consider two states ϱ and σ of an N -party system described by the Hilbert space $\mathcal{H}_{A_1 \dots A_N}$. The question of local unitary equivalence can be mathematically formulated in the following way: *Does there exist a unitary transformation $U = U_1 \otimes \dots \otimes U_N$, such that $\varrho = U\sigma U^\dagger$?*

In order to study this problem of local unitary equivalence we firstly, simplify our task. Specifically, we divide the total Hilbert space in the following way

$$\mathcal{H}_{A_1 \dots A_N} = \mathcal{H}_{A_1} \otimes \dots \otimes \mathcal{H}_{A_N} = \mathcal{H}_X \otimes \mathcal{H}_Y. \quad (2.1)$$

where X and Y are two sets of subsystems A_k . If we would have some criteria for bi-partite systems being locally unitary equivalent, then we can decide whether the following is true

$$\varrho = U_X \otimes U_Y \sigma U_X^\dagger \otimes U_Y^\dagger. \quad (2.2)$$

If the answer is “no”, then the states are not locally unitary equivalent. If the answer is “yes”, then we continue our investigation by checking whether the pair of states

$$\begin{aligned} \varrho_X &= \text{Tr}_Y \varrho \quad \text{and} \quad \sigma_X = \text{Tr}_Y \sigma \\ \varrho_Y &= \text{Tr}_X \varrho \quad \text{and} \quad \sigma_Y = \text{Tr}_X \sigma \end{aligned} \quad (2.3)$$

are locally unitary equivalent. That is, we again, divide each of the subsystems X, Y into two smaller sets and we use the criteria for bi-partite system. Using this method recursively we finally partition the multi-partite system into its components A_k and give an answer to the question of locally unitary equivalence. From here it follows that it is enough to have a criterion for the bi-partite composite systems with the help of which multi-partite systems can be studied. In what follows, we present some necessary conditions for states to be locally unitary equivalent.

A. Invariants of local unitaries

1. **Eigenvalues.** It is well known that the necessary and sufficient condition for two states to be unitarily equivalent is that they have the same set of eigenvalues. Let us denote by $\text{eig}(\varrho)$ the set of eigenvalues of the density matrix ϱ . Then

$$\varrho' = U \varrho U^\dagger, \quad \text{iff} \quad \text{eig}(\varrho') \equiv \text{eig}(\varrho). \quad (2.4)$$

2. **Eigenvalues of the reduced states.** The reduced density matrices ϱ_X and ϱ_Y change after the action of the local unitary transformation $U_X \otimes U_Y$ as follows: for all U_X and U_Y

$$\begin{aligned} \varrho'_X &= \text{Tr}_Y [U_X \otimes U_Y \varrho U_X^\dagger \otimes U_Y^\dagger] = U_X \varrho_X U_X^\dagger, \\ \varrho'_Y &= \text{Tr}_X [U_X \otimes U_Y \varrho U_X^\dagger \otimes U_Y^\dagger] = U_Y \varrho_Y U_Y^\dagger. \end{aligned} \quad (2.5)$$

Now we can define the first invariant: local unitary transformations do not change the eigenvalues of the reduced density matrices, which describe states of the subsystems.

For any state ϱ' such that $\varrho' = U_X \otimes U_Y \varrho U_X^\dagger \otimes U_Y^\dagger$ it is valid that

$$\begin{aligned} \text{eig}(\varrho_X) &\equiv \text{eig}(\varrho'_X), \\ \text{eig}(\varrho_Y) &\equiv \text{eig}(\varrho'_Y). \end{aligned} \quad (2.6)$$

We note that this relation is valid for any operator ϱ , not only for density operators.

3. Eigenvalues of the reduced eigenprojectors.

Quantum states can be expressed in a spectral form. Specifically, let $U = U_X \otimes U_Y$ and $\varrho = \sum_k \lambda_k P_k$, then $\varrho' = U \varrho U^\dagger = \sum_k \lambda_k P'_k$, where P_k and $P'_k = U P_k U^\dagger$ are eigenprojectors belonging to the eigenvalue λ_k . Since the unitary transformation under consideration is local, i.e. $U = U_X \otimes U_Y$, we obtain the implication

$$\begin{aligned} \varrho' = U \varrho U^\dagger \quad \text{then} \quad \dim(P_k) &= \dim(P'_k) \\ \text{eig}(\text{Tr}_Y P_k) &\equiv \text{eig}(\text{Tr}_Y P'_k) \\ \text{eig}(\text{Tr}_X P_k) &\equiv \text{eig}(\text{Tr}_X P'_k), \end{aligned} \quad (2.7)$$

where $\dim(P_k)$ denotes the dimension of the projection P_k defined as the number of its nonzero eigenvalues.

4. **Partial transposition.** We use the following identity for any operators K_X, L_Y, M_X, N_Y

$$(K_X \otimes L_Y \varrho M_X \otimes N_Y)^{T_Y} = K_X \otimes N_Y^T \varrho^{T_Y} M_X \otimes L_Y^T \quad (2.8)$$

where by T_Y we denote the partial transposition of the subsystem Y . A similar identity is fulfilled also for the partial transposition, T_X , of the system X . Note, that the partial transposition depends on the choice of the local basis. Introducing the local unitary transformation $U_X \otimes U_Y$ we obtain

$$\begin{aligned} (\varrho')^{T_Y} &= (U_X \otimes U_Y \varrho U_X^\dagger \otimes U_Y^\dagger)^{T_Y} \\ &= U_X \otimes (U_Y^\dagger)^T \varrho^{T_Y} U_X^\dagger \otimes U_Y^T \\ &= W \varrho^{T_Y} W^\dagger \end{aligned} \quad (2.9)$$

where $W = U_X \otimes (U_Y^T)^\dagger$ is a unitary operator. Its unitarity follows from the fact that the transposed unitary operator is a unitary operator. The unitary transformation leaves the eigenvalues of the operator ϱ^{T_Y} unchanged. Therefore, we obtain the second invariant associated with density matrices which are invariant under local unitary transformations. Namely, if $\varrho' = U_X \otimes U_Y \varrho U_X^\dagger \otimes U_Y^\dagger$ then

$$\begin{aligned} \text{eig}(\varrho^{T_Y}) &\equiv \text{eig}(\varrho'^{T_Y}) \\ \text{eig}(\varrho^{T_X}) &\equiv \text{eig}(\varrho'^{T_X}). \end{aligned} \quad (2.10)$$

B. Local unitary equivalence for pure states

Let us consider two pure states $|\psi\rangle$ and $|\phi\rangle$ of a bi-partite system written in the Schmidt-decomposed form

$$\begin{aligned} |\psi\rangle &= \sum_k \sqrt{\lambda_k} |k\rangle_X \otimes |k\rangle_Y \\ |\phi\rangle &= \sum_k \sqrt{\mu_k} |k'\rangle_X \otimes |k'\rangle_Y. \end{aligned} \quad (2.11)$$

From the equality $\lambda_k = \mu_k$ (for all k) the validity of the second invariant property follows. That is the reduced states have the same set of eigenvalues, namely $\{\lambda_k\}$. The first invariant property is fulfilled trivially for pure states. Moreover, let us define local unitaries by the equations $U_X |k\rangle_X = |k'\rangle_X$ and $U_Y |k\rangle_Y = |k'\rangle_Y$. Then the states (2.11) are locally unitary equivalent and $|\phi\rangle = U_X \otimes U_Y |\psi\rangle$.

Theorem 1

Let us consider two pure states $|\psi\rangle$ and $|\phi\rangle$ of a bi-partite system. These state are locally unitary equivalent if and only if their coefficients in the Schmidt decomposition are equal, i.e. the sets of eigenvalues of reduced states coincide.

Example 1.

This simple theorem can be rather useful. In particular, we can use it to prove that any GHZ state is locally equivalent to one EPR pair and some ancilla system in an arbitrary state. It means we are able to prepare a GHZ state from an EPR pair a vice versa [1]. Let us define two states of N qubits $|GHZ\rangle_{1\dots N} = (|0^{\otimes N}\rangle + |1^{\otimes N}\rangle)/\sqrt{2}$ and $|\Psi\rangle_{1\dots N} = |EPR\rangle_{12} \otimes |\phi\rangle_{3\dots N}$, where $|EPR\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$. Since $\text{Tr}_{2\dots N} |GHZ\rangle\langle GHZ| = \text{Tr}_{2\dots N} |\Psi\rangle\langle\Psi| = \frac{1}{2}\mathbb{1}$ we know that the eigenvalues of the reduced states coincide. Specifically, let us consider a splitting of the whole system into a single qubit denoted as “1” and the rest of the system. It means that these states are locally (with respect to the given splitting) unitary equivalent.

Example 2.

Let us consider a state (for more details see [7])

$$|X\rangle = \frac{1}{\sqrt{2^n}} \sum_{k=0}^{2^n-1} |k\rangle_X \otimes |\alpha_k\rangle_Y \quad (2.12)$$

of $N = 2n + 1$ qubits located on a ring. This system is divided into two subsystem of n neighboring qubits (subsystem X) and $(n + 1)$ neighboring qubits (subsystem Y). In this notation $|k\rangle_X$ represents the state of n qubits, if k is written in a binary form. The state $|\alpha_k\rangle_Y = \sum_l (-1)^{d_k + d_l + d_{k,l}} |l\rangle_Y$ where $|l\rangle_Y$ is the state of the rest of the $n + 1$ qubits, if l is written in a binary form. The number k and l define the state of the ring in the sense of its binary expression. From here we can determine the state of each individual qubit. The numbers k give us the qubits in the state $|1\rangle$ and we can write $\{i, \dots, j\}$ to represents the position of the

qubits in this state. Similarly, l determines the set $\{r, \dots, s\}$. We define $d_k = d_{\{i, \dots, j\}} := \sum_{i < j} d(i, j)$, $d_{k,l} = \sum_{a \in \{i, \dots, j\}, b \in \{r, \dots, s\}} d(a, b)$, where $d(i, j)$ is the shortest distance between two qubits (in the state $|1\rangle$) on a ring (i.e. number of steps needed to get from qubit in position i into the position j around the ring, or the smallest number of qubits lying between the i -th and j -th position). The state (2.12) is one of the eigenstates of the Ising model (for more details see [7]).

For the purpose of our example it is important that Eq. (2.12) represents the Schmidt form of the state $|X\rangle$ with all Schmidt coefficients equal to $1/\sqrt{2^n}$ and the Schmidt number (i.e. number of nonzero Schmidt coefficients) is equal to 2^n . Let us define a state

$$|\Psi\rangle = |\phi\rangle_{Y_{n+1}} \otimes \left(\bigotimes_{j=1}^n |EPR\rangle_{X_j Y_j} \right), \quad (2.13)$$

where $|EPR\rangle_{X_j Y_j} = (|0\rangle_{X_j} \otimes |0\rangle_{Y_j} + |1\rangle_{X_j} \otimes |1\rangle_{Y_j})/\sqrt{2}$ is the EPR pair shared between the subsystems X and Y , such that j represents the position of the qubit in this subsystems. We can express the state (2.13) in the Schmidt form

$$|\Psi\rangle = \frac{1}{\sqrt{2^n}} \sum_{k=0}^{2^n-1} |k\rangle_X \otimes |\beta_k\rangle_Y, \quad (2.14)$$

where $|\beta_k\rangle_Y = |k\rangle_{Y'} \otimes |\phi\rangle_{n+1}$ and Y' denotes the subsystem Y without one of its qubit at the $(n + 1)$ -th position. Since the set of Schmidt coefficients for these states coincides, we conclude that the states (2.12) and (2.13) are locally unitarily equivalent. We stress that this equivalence is with respect to the splitting of the ring into two parts X and Y . Therefore, from the point of view of entanglement between systems X and Y these states are equivalent, and we are able to prepare n EPR pairs between these systems by a local transformation $U = \mathbb{1}_X \otimes U_Y$, where U_Y is determined by the equality $U_Y |\alpha_k\rangle_Y = |\beta_k\rangle_Y$. Finally we note, that this transformation is not uniquely defined, because of the degeneracy of Schmidt coefficients.

C. Counterexample for mixed states

Unfortunately, Theorem 1 is valid only for pure states and cannot be generalized for arbitrary impure states. In what follows we will show a simple counterexample. The idea behind this example is that we cannot create entanglement by a local unitary transformation and that the spectral decomposition of a density matrix is unique. It means the separable pure state in the spectral decomposition cannot evolve into entangled one.

Example 3.

Let us consider two states ϱ_{XY} and σ_{XY} of a bi-partite system $\mathcal{H}_X \otimes \mathcal{H}_Y$, where $\dim \mathcal{H}_Y = 2$ and $\dim \mathcal{H}_X = 8$.

This means the system of four qubits is divided into a one-qubit and a three-qubit subsystems. Let the set of eigenvalues be the same

$$\text{eig}(\varrho_{XY}) = \text{eig}(\sigma_{XY}) = \left\{ \frac{1}{4}, \frac{3}{8}, \frac{5}{16}, \frac{1}{16} \right\} \quad (2.15)$$

and the corresponding eigenvectors be

$$\begin{pmatrix} \frac{1}{\sqrt{4}}(|0,1\rangle + |0,2\rangle + |0,4\rangle + |1,0\rangle) \\ |1,7\rangle \\ |0,5\rangle \\ \frac{1}{\sqrt{3}}(|1,1\rangle + |1,2\rangle + |1,4\rangle) \end{pmatrix} \quad (2.16)$$

for ϱ_{XY} and

$$\begin{pmatrix} |1,7\rangle \\ \frac{1}{\sqrt{3}}(|0,7\rangle + |1,3\rangle + |0,5\rangle) \\ |0,0\rangle \\ |0,1\rangle \end{pmatrix} \quad (2.17)$$

for σ_{XY} . It is easy to check that these two states also have the same sets of eigenvalues of their reduced density matrices, i.e.

$$\begin{aligned} \text{eig}(\varrho_X) &= \text{eig}(\sigma_X) = \left\{ \frac{1}{2}, \frac{1}{2} \right\} \\ \text{eig}(\varrho_Y) &= \text{eig}(\sigma_Y) = \left\{ \frac{3}{8}, \frac{5}{16}, \frac{1}{4}, \frac{1}{16} \right\} \end{aligned} \quad (2.18)$$

So for the state under considerations the conditions of Theorem 1 are satisfied. On the other hand, in contradiction to the general properties of local unitaries, we see the violation of the property the entanglement creation prohibited by local unitary transformation. In other words, the condition of the same set of reduced eigenvalues of eigenprojectors (invariant A3) is not fulfilled.

III. LOCAL UNITARY EQUIVALENCE FOR THE MIXED STATES

We have shown that Theorem 1 is not valid for all states. Certainly, it can be trivially generalized to a class of the so called *factorizable states*, given by density operators $\varrho_{XY} = \varrho_X \otimes \varrho_Y$. On the other hand, it is not clear whether the invariants presented in Section II provide us with a definite answer for the problem of the local unitary equivalence for mixed states. To clarify the situation we formulate the following theorem

Theorem 2

Suppose two non-degenerate states ϱ and σ of the composite system $X + Y$ are given, i.e.

$$\varrho = \sum_k \lambda_k |\psi_k\rangle \langle \psi_k|, \quad \sigma = \sum_k \mu_k |\phi_k\rangle \langle \phi_k|. \quad (3.1)$$

and $\lambda_k \neq \lambda_l, \mu_k \neq \mu_l$ for $k \neq l$. Let us express the vectors $|\psi_1\rangle$ and $|\phi_1\rangle$ in their Schmidt bases and fix these two basis on $\mathcal{H}_X \otimes \mathcal{H}_Y$. If for each k , $\lambda_k = \mu_k$ and the coefficients of the corresponding eigenvectors $|\psi_k\rangle = \sum_{mn} \alpha_{mn}^k |mn\rangle$ and $|\phi_k\rangle = \sum_{mn} \beta_{mn}^k |m'n'\rangle$ (where $|mn\rangle$ and $|m'n'\rangle$ are the previously mentioned fixed bases) coincide, i.e. $\alpha_{mn}^k = \beta_{mn}^k$, then the states ϱ, σ are locally unitary equivalent. The local transformation is given by $U_X|m\rangle = |m'\rangle$ and $U_Y|n\rangle = |n'\rangle$.

From the construction it is clear that the theorem is valid. Note that the choice of the fixed basis is given by the Schmidt expression of one of the eigenvectors, but the result doesn't depend on it. In the degenerate case we cannot use the argument with the Schmidt decomposition, because then the eigenprojectors are no longer one dimensional.

We conclude this section by saying that in general by using invariants of local unitaries we are able only to determine whether the states under consideration are not locally unitary equivalent.

IV. LOCAL CONNECTION OF MULTI-PARTITE AND BIPARTITE ENTANGLEMENT

In this section we generalize Example 2 of Section III. Specifically, let us consider a pure state $|\psi\rangle_{A_1, \dots, A_N}$ of a composite system consisting of N qudits. Let us split this system into two multi-partite subsystems X and Y . By calculating the entropy of one of the subsystems we can determine the degree of entanglement between these two parties. An interesting question is whether with the help of a local transformation with respect to the splitting we are able to prepare a state where the entanglement is shared only between the specific pairs of qudits such that one particle of each pair belongs to the system X while the second belongs to the system Y . In other words, the qudits belonging to the subsystem X (Y) are mutually separable. It means that the multi-partite entanglement shared between the subsystems X and Y can be arranged by bi-partite entanglement shared between the pairs of qudits just by performing an appropriate local unitary transformation $U = U_X \otimes U_Y$ (see Fig. 1).

Let us denote by $A_1, \dots, A_N$ the qudits of the whole system and by $X_1, \dots, X_n$ ($Y_1, \dots, Y_m$) the qudits in a subsystem X (Y). Note that $n + m = N$. We also introduce a function $E_{XY}(|\psi\rangle_{XY})$ which describes the degree of entanglement shared between the systems X and Y . Suppose $n \leq m$. The most general pure state of the whole system, for which the entanglement between the subsystems X and Y is equal to the entanglement shared by pairs of qudits is given by the expression

$$|\Phi\rangle_{XY} = \left(\bigotimes_{j=1}^n |\chi\rangle_j \right) \otimes |rest\rangle_{Y_{n+1}, \dots, Y_m} \quad (4.1)$$

where $|\chi\rangle_j = \sum_k \sqrt{a_{kj}} |\chi_k\rangle_{X_j} \otimes |\chi_k\rangle_{Y_j}$ is state written in Schmidt basis. The entanglement between the systems

X and Y which are in the state (4.1) is given can be represented by the expression

$$E_{XY}(|\Phi\rangle_{XY}) = \sum_{jk} a_{k_j} \log a_{k_j}. \quad (4.2)$$

The equality $E_{XY}(|\psi\rangle_{XY}) = E_{XY}(|\Phi\rangle_{XY})$ is not sufficient to determine whether the states $|\psi\rangle_{XY}$ and $|\Phi\rangle_{XY}$ are locally unitary equivalent. Let us rewrite both states in their Schmidt bases

$$\begin{aligned} |\psi\rangle_{XY} &= \sum_{r=0}^{d^n-1} \sqrt{q_r} |\alpha_r\rangle_X \otimes |\beta_r\rangle_Y, \\ |\Phi\rangle_{XY} &= \sum_{r=0}^{d^n-1} \sqrt{p_r} |r\rangle_X \otimes (|r\rangle_{Y_1, \dots, Y_n} \otimes |rest\rangle_{Y_{n+1}, \dots, Y_m}), \end{aligned} \quad (4.3)$$

where $d = \dim(\mathcal{H}_{A_j})$, r is a base d number which represents the state of n qudits, e.g. $r = 4$ in a binary form represents a state of 5 qubits $|0, 0, 1, 0, 0\rangle$, or in the base 3 form it represents a state of 5 qutrits $|0, 0, 0, 1, 2\rangle$.




FIG. 1. Schematic representation of a connection between multi-partite and bipartite entanglement via local operations. On the left side of the figure the double line represents the shared multi-partite entanglement between X and Y . Each line in the right side of the figure represents an entangled pair of qudits $|\chi\rangle_j$, i.e. bipartite entanglement.

The eigenvalues of the reduced states are $\{q_r\}$ for the state $|\psi\rangle_{XY}$ and $\{p_r = a_{k_1} \dots a_{k_n}\}$ for the state $|\Phi\rangle_{XY}$, where $r = (k_1, \dots, k_n)$ in the number base- d form, i.e. $k_j = 0, 1, \dots, d-1$. Theorem 1 implies that these sets of eigenvalues have to be identical in order to ensure local unitary equivalence. Our aim is to find some condition under which the equations

$$q_r = p_r = p_{(k_1, \dots, k_n)} = a_{k_1} \dots a_{k_n}. \quad (4.4)$$

are satisfied. In what follows we have d^n equations with nd undetermined variables. These equation, in general, do not have a solution. In that case the states are not locally unitary equivalent and the entanglement shared by systems X and Y cannot be prepared by local unitaries from bi-partite entanglement shared between the subsystems. If the system of equations (4.4) has a solution then the local unitary transformations are given by relations

$$\begin{aligned} U_X |\alpha_r\rangle_X &:= |\alpha_r\rangle_X \\ U_Y |\beta_r\rangle_Y &:= |r\rangle_{Y_1, \dots, Y_n} \otimes |rest\rangle_{Y_{n+1}, \dots, Y_m}. \end{aligned} \quad (4.5)$$

We note, that from above, the local unitary equivalence between the $|X\rangle$ state and n EPR pairs shared between systems X and Y follows (see Section II).

V. CONCLUSION

In this paper we have analyzed the problem of finding under which conditions two states ϱ and σ of a multi-partite system are locally unitary equivalent. We have presented a set of conditions which have to be satisfied in order that the two states are locally unitary equivalent. We have presented a set of parameters which play the role of invariants under the action of local unitary transformations. In addition, we have shown how these invariants can be used to characterize the local unitary equivalence of states.

We have shown that the problem of local unitary equivalence for multi-partite systems can be reduced to the problem of analysis of bi-partite systems. Specifically, it is enough to investigate the case of bi-partite local unitary equivalence and to show sufficient conditions of this property for non-degenerate density matrices.

We left two questions opened: the local unitary equivalence for degenerate density matrices and the characterization of the connection between multi-partite and bipartite entanglement, since, as it is well known that there exists an intrinsic multi-partite entanglement [8,9].

Finally, we have analyzed multi-qudit systems. We have shown, that when analyzing local unitary equivalence of two states it is useful to consider the splitting of the whole system into two subsystems. X and Y . In this case a system of equations of the form (4.4) results. We have shown that these equations might not have a solution from which it directly follows that the state under consideration cannot be prepared by local unitary transformations from any number of partially entangled pairs shared between two systems X and Y (i.e. these states cannot be prepared just using a bi-partite entanglement as a resource and local transformations on subsystems X and Y).

ACKNOWLEDGMENTS

We thank Mark Hillery for discussions and for reading the manuscript. V.B. thanks Chris Fuchs for a stimulating discussion. This work was supported by the IST project EQUIP under the contract IST-1999-11053 and by the grant GACR 201/01/0413.

- [1] See for example: M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [2] J. Gruska, *Quantum Computing* (McGraw-Hill, London, 1999).
- [3] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer Academic Publishers, 1995).
- [4] K.G.H.Vollbrecht,R.F.Werner,*Entanglement measures under symmetry*, LANL preprint archive [quant-ph/0010095](#)
- [5] A.Sudbery, *On local invariants of pure three-qubit states* *J.Phys.A* **34**, 643-652 (2001), also available at LANL preprint archive [quant-ps/0001116](#)
- [6] H.Barnum, N.Linden, *Monotones and invariants for multi-particle quantum states*, LANL preprint archive [quant-ph/0103155](#)
- [7] P.Štelmachovič and V.Bužek, *Entanglement in Ising model* (in preparation).
- [8] A. V. Thapliyal, Phys. Rev. A **59**, 3336 (1999); J. Kempe, Phys. Rev. A **60**, 910 (1999).
- [9] S. Hill and W. K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997); W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998). V. Coffman, J. Kundu, and W. K. Wootters, [quant-ph/9907047](#) (1999) W. K. Wootters, [arXiv quant-ph/0001114](#) (2000).